



FUNGICIDE

TRANSOM[®] 4.5 F

Liquid Flowable Systemic Turf and Ornamental Fungicide

For control of a broad spectrum of diseases of bedding, flowering, herbaceous and tropical foliage plants, shrubs, trees and flowers, containerized woody shrubs and trees and turfgrass.

ACTIVE INGREDIENT:

Thiophanate-methyl	46.2%
OTHER INGREDIENTS:	53.8%
TOTAL:	100.0%

(Contains 4.5 Pounds Thiophanate-methyl Per Gallon)

EPA Reg. No. 72112-2

EPA Est. No. indicated by the first two letters of the batch number on this package.
(AL) 228-IL-002, (YU) 67545-AZ-001

Product of China or India.
Formulated in the United States with U.S. and imported ingredients.



8 95217 00112 0

Net Contents:

2.5 Gallon (9.46 liters)

For PRODUCT USE Information
Call 1-770-619-9832.

Manufactured for:
PROKÖZ, Inc.
1720 Windward Concourse
Suite 150
Alpharetta, GA 30005

**KEEP OUT
OF REACH
OF CHILDREN
CAUTION**

For FIRST AID and
PRECAUTIONARY STATEMENTS:
See Inside Booklet.

For MEDICAL and TRANSPORTATION
Emergencies ONLY
Call 24 Hours A Day 1-800-424-9300.

FIRST AID

If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Harmful if swallowed or inhaled. Harmful if absorbed through skin. Avoid breathing spray mist. Avoid contact with eyes, skin or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, polyvinyl chloride (PVC) \geq 14 mils or viton \geq 14 mils. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

Handlers mixing, loading and applying the product as a dip must wear:

- Coveralls over long-sleeved shirt and long pants;
- Chemical-resistant gloves;
- Chemical-resistant footwear plus socks;
- A chemical-resistant apron.

All other mixers and loaders and applicators must wear:

- Long-sleeved shirt and long pants;
- Shoes plus socks;
- Chemical-resistant gloves for all mixers and loaders and for applicators using hand-held equipment, and
- Chemical-resistant apron for mixers, loaders and other handlers exposed to the concentrate.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Engineering Controls

When handlers use enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Not for homeowner use. For use by individuals/firms licensed or registered by the state to apply ornamental or turf pest control products. Do not apply with fixed wing or rotary aircraft.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exemption: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water), is:

- Coveralls over long-sleeved shirt and long pants,
- Chemical-resistant gloves made of any waterproof material,
- Chemical-resistant footwear plus socks,
- Chemical-resistant headgear for overhead exposures.

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter treated areas without protective clothing until spray has dried.

GENERAL INFORMATION

TRANSOM® 4.5 F provides broad spectrum disease control on containerized woody, flowering, herbaceous, and tropical foliage ornamental plants and trees, and turfgrasses.

Add the required amount of TRANSOM® 4.5 F after shaking well to a partially-filled tank agitated by mechanical or hydraulic means then add the remaining required amount of water. Maintain continuous agitation during application to keep the material in suspension. Make fresh daily.

Make applications of TRANSOM® 4.5 F with ground equipment, using sufficient spray volume to provide thorough coverage. Do not tank mix TRANSOM® 4.5 F with copper containing materials or with highly alkaline pesticides such as Bordeaux mixture or lime sulfur. No claim of compatibility with other pesticides is implied. Under conditions of severe disease pressure or when application intervals are shorter than 14 days due to persistent rainfall, use the higher concentration or rates provided in this label. Contact your local State Extension Service specialist for application schedule recommendations.

Important: If recommended treatments of TRANSOM® 4.5 F are ineffective, a tolerant strain of fungus may be present. Consult your PROK6Z representative or distributor, your local State Agricultural Experiment Station or State Agricultural Extension Service for advice on prompt use of some other labeled fungicide.

CHEMIGATION INSTRUCTIONS

For Ornamental Use Only

Apply this product only through pressurized drench (flood), sprinkler, or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation systems.

Do not connect chemigation system to a public water system.

Crop injury, lack of effectiveness, or illegal residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, contact your State Extension Service specialists, equipment manufacturers, or other experts.

Do not connect any irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

Should the need arise, a person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments.

Pressurized Drench (Flood) System

Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump, (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Systems using a gravity-flow pesticide-dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops.

Sprinkler Chemigation

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump, (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.

Drip (Trickle) Chemigation

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

HORTICULTURAL APPLICATIONS

GREENHOUSE, NURSERY AND LANDSCAPE

TRANSOM® 4.5 F provides broad spectrum control of many foliar, stem, and below ground diseases on a wide range of horticultural plants grown or maintained under a variety of conditions. Apply TRANSOM® 4.5 F 14-21 days prior to when a particular disease usually appears and at the very latest, upon first sign of disease. Spray intervals usually range from 7 to 14 days with 14 days being for preventative treatments and the 7-day interval for times when conditions are judged acceptable for disease development. For hard-to-wet foliage, an acceptable wetting agent may be added to the spray tank to increase product efficacy. Use of a spreader-sticker is recommended when excessive and repeated foliar wetting occurs. TRANSOM® 4.5 F may be used to control listed diseases on non-commercial fruit and nut trees. Do not use fruit or nuts from trees treated with this product as food. Do not apply this product to home orchards or backyard fruit trees after fruit set.

Note: TRANSOM® 4.5 F has been determined to be safe for use on the plant types listed in these directions for use based on cumulative data derived from research product trials and historical field use. As all species and cultivars have not been tested, it is recommended that trial applications be performed if a user wishes to make an application to a plant type not listed on the label but found on a similar use site and for a disease that is listed on the label. To conduct a trial application, apply at least two applications to at least 25 trial plants at the highest concentration, 7 days apart. Evaluate 7 days after the last application before initiating full-scale application. Do not use this product on the following plants: Swedish Ivy (*Neprolepsis exaltata*), Boston Fern (*Plectranthus australis*), and Easter Cactus (*Hatiora gaertneri*).

Plant Type	Such as but not limited to:
Herbaceous Bedding	Ageratum, Begonia, Canna, Coleus, Dahlia, Dusty Miller, Foxglove, Fuchsia, Geranium, Impatiens, Lavender, Marigold, Pansy, Petunia, Pinks, Primrose, Salvia, Statice, Strawflower, Tickseed, Verbena
Flowering	Chrysanthemum, Hydrangea, Hollyhock, Iris, Lily, Poinsettia
Tropical Foliage	Dieffenbachia, Dracaena, English Ivy, Philodendron, Pothos
Woody Ornamentals	Azalea, Hibiscus, Holly, Ligustrum, Rhododendron, Rose, Pyracantha
Evergreen Trees	Douglas Fir, Fir, Larch, Pine, Spruce
Deciduous Trees*	Ash, London Plane, Maple, Oak, Sycamore, Walnut
Flowering Trees*	Cherry, Crabapple, Hawthorn, Mountain Ash, Pear

*Do not use fruit or nuts from treated trees as food or feed.

FOLIAR SPRAY APPLICATIONS

Hydraulic Application Mixing Instructions

After shaking the product container, add the required amount of TRANSOM® 4.5 F to a partially-filled spray tank agitated by mechanical or hydraulic means and then add the remaining recommended volume of water. Maintain continuous agitation to keep the material in suspension and apply with properly calibrated spray equipment.

Application Concentrations (Mechanical or Hand-Held):

Use the recommended amount of TRANSOM® 4.5 F per 100 gallons of water for the prevention and control of the diseases shown below.

FOLIAR DISEASES

Disease(s) Controlled	Concentration of TRANSOM® 4.5 F fl. oz./100 gals.	Remarks
Anthracoese <i>Colletotrichum</i>	10.75 – 20	Apply as buds break or at first sign of disease. Repeat application at 7- to 14-day intervals as needed during disease period.
Black Spot of Rose <i>Diplocarpon rosae</i>	10.75 – 20	Apply early summer or at first sign of disease. Repeat application every 7-14 days as needed during disease period.
Brown Rot and Blight <i>Monilinia, Sclerotini, Whetzelinia</i>	10.75 – 20	Apply late spring or at first sign of disease. Repeat application every 7-14 days as needed during the disease period.
Fusicladium and Venturia Leaf Scabs on: Crabapple, Hawthorn, Pear, Mountain Ash, Pyracantha, etc.	10.75 – 20	Apply as buds break. Repeat application every 7-14 days during disease period. Effective control requires coverage during expansion. Rotations with chlorothalonil or propiconazole can be utilized.

(continued on next panel)

FOLIAR DISEASES (continued)

Disease(s) Controlled	Concentration of TRANSOM® 4.5 F fl. oz./100 gals.	Remarks
Leaf Spots and Blights caused by: <i>Ascochyta</i> , <i>Blumeriella</i> , <i>Botrytis</i> , <i>Cercospora</i> , <i>Coccomyces</i> , <i>Corynespora</i> , <i>Curvularia</i> , <i>Didymellina</i> , <i>Entomosporium</i> , <i>Fabraea</i> , <i>Fusarium</i> , <i>Ramularia</i> , <i>Rhizoctonia</i> , <i>Marssoninia</i> , <i>Mycosphaerella</i> , <i>Myrothecium</i> , <i>Phoma</i> , <i>Physalospora</i> , <i>Schizothyrium</i> , <i>Septoria</i> , <i>Sphaceloma</i>	10 – 14.5	Make applications when disease symptoms first appear. Repeat every 7-14 days during disease period. Rotations with chlorothalonil may be used.
Ovulinia Blight	7.25 – 20	Apply as flowers open. Repeat every 7-14 days during disease period.
Powdery Mildews <i>Erysiphe</i> , <i>Microsphaera</i> , <i>Phyllactinia</i> , <i>Podosphaera</i> , <i>Oidium</i> , <i>Sphaerotheca</i>	10 – 20	Apply when disease first appears and repeat application every 7-14 days. Rotations with other effective products is recommended.
Rust Diseases caused by: <i>Puccinia</i> , <i>Gymnosporangium</i> , <i>Uromyces</i>	10.75 – 20	Apply late spring or when symptoms first appear. Repeat application every 7-14 days during disease period. Rotations with other effective products is recommended.
Tip Blight of Pine <i>Sphaeropsis sapinea</i> , <i>Diplodia pinea</i>	14.5 – 20	Begin applications in the spring when new growth starts. Make a second application just prior to needle emergence from the sheath and a third application 7 days later. Ensure thorough coverage.
Twig Blights, Cankers, and Diebacks <i>Diaporthe</i> , <i>Kabatina</i> , <i>Phoma</i> , <i>Phomopsis</i>	14.5 – 20	Apply when symptoms first appear. Repeat application every 7-14 days during disease period.

Adjuvants: Where rainfall and/or overhead irrigation is the norm, use of a compatible spreader/sticker is suggested. Where wetting of foliage is difficult, use a compatible wetting agent. Follow the phytotoxicity precautions described in the "HORTICULTURAL APPLICATIONS" section of this label.

SOIL DRENCH APPLICATIONS

Mixing Instructions: Add the required amount of TRANSOM® 4.5 F to a partially-filled tank agitated by mechanical or hydraulic means. Add the remaining required amount of water. Maintain continuous agitation throughout application to keep the material in suspension.

Application Concentrations/Rates and Timing for Disease Control: Use 7.5 to 20 fl. oz. of TRANSOM® 4.5 F per 100 gallons of water. Apply as a drench or heavy spray at the rate of 1/2 to 2 pints per square foot (100 gallons per 400 to 1,600 square feet). For small pots and shallow flats up to 4 inches in size, apply at 1 pint per square foot. For containers and pots 4 inches or larger, refer to the following table for the volume to apply. Repeat applications may be made at 4- to 8-week intervals depending on disease presence and conditions for disease development.

Container Type	Volume to Apply/Container	
	1 pt./sq. ft. Rate	2 pt./sq. ft. Rate
4 inch	2 fl. oz.	
5 inch	2-1/2 fl. oz.	
6 inch		6-1/2 fl. oz.
7 inch		8-1/2 fl. oz.
8 inch		11 fl. oz.
9 inch		14 fl. oz.
10 inch		17-1/2 fl. oz.

For containers larger than 10 inches, a drench volume of 2-1/2 to 3 pints per square foot of surface area may be required.

Plant Types: Containerized woody shrubs, trees, herbaceous/bedding, flowering, and tropical foliage plants and flowers and bedding plants in the landscape.

Note: Do not apply this product to plug trays or seedling flats at time of seeding.

Soil Diseases Controlled: *Botrytis*, *Cylindrocladium*, *Fusarium*, *Gliocladium*, *Myrothecium*, *Penicillium*, *Ramularia*, *Rhizoctonia*, *Sclerotinia* and *Thielaviopsis*.

Note: Pythium, Phytophthora and Cylindrocladium spathiphylli are not controlled by TRANSOM® 4.5 F.

PLANT DIP TREATMENT

Mixing Instructions: Mix as described in the "FOLIAR" and "SOIL DRENCH APPLICATIONS" sections of this label. Maintain continuous agitation during application.

Application Concentration and Dipping Time

Plants or Cuttings: Use 14.5 to 21.5 fl. oz. of TRANSOM® 4.5 F per 100 gallons of water. Immerse plants or cuttings for 10 to 15 minutes, remove and allow to drain and dry. Wear protective clothing as described under the "PERSONAL PROTECTIVE EQUIPMENT" section of this label.

Bulbs, Corms, Tubers and Rhizomes: Use 14.5 to 33 fl. oz. of TRANSOM® 4.5 F per 100 gallons of water or 2 teaspoons of TRANSOM® 4.5 F per gallon of water. Soak cleaned bulbs for 15 to 30 minutes in warm dip (80-85°F) within 48 hours of digging. Dry bulbs after treatment. If bulbs are for forcing, treat bulbs that have been heat-cured.

Plant Types: Plants, cuttings, cane sections of woody herbaceous, flowering and tropical foliage plants. Bulbs, corms, tubers, and rhizomes of plants such as but not limited to Caladium, Easter Lily, Tulip, Gladiolus, Daffodil, Iris.

Diseases Controlled: *Botrytis*, *Cylindrocladium*, *Fusarium*, *Gliocladium*, *Myrothecium*, *Penicillium*, *Ramularia*, *Rhizoctonia*, *Sclerotinia*, *Thielaviopsis*.

TURF APPLICATIONS

Not for homeowner use. For use only by certified applicators or those under their immediate supervision. Do not apply with fixed wing or rotary aircraft. Not for use on turf being grown for sale or other commercial use as sod.

TRANSOM® 4.5 F may be used against certain foliar and soil diseases for use on all turf types such as golf course greens, tees and fairways, athletic fields, cemeteries, parks, and commercial and home lawns. TRANSOM® 4.5 F may be used both preventively and curatively and is not phytotoxic. Do not use TRANSOM® 4.5 F on turf being grown for sale or other commercial uses as sod.

Mixing Instructions: Add the required amount of TRANSOM® 4.5 F to a partially-filled tank agitated by mechanical or hydraulic means. Add the remaining required amount of water. Maintain continuous agitation to keep the material in suspension. For best results, use spray mixture the same day it is prepared.

Turf Types: All cool-season and warm-season grasses (such as but not limited to Bentgrasses, Bermudagrasses, Bluegrasses, Fescues, Ryegrasses, St. Augustinegrasses and Zoysiagrasses) or their mixtures.

Disease(s) Controlled	Rate of TRANSOM® 4.5 F fl. oz./1,000 sq. ft.**	Remarks
Anthracnose <i>Collectotricum graminicola</i>	2 – 3-1/2 (3-1/2 – 5-1/3)*	Apply when disease first appears. Make additional applications at 14-day intervals as needed. Allow spray to dry on leaves with no “watering-in”.
Dollar Spot <i>Sclerotinia homoeocarpa</i> Copper Spot <i>Gloeocerospora sorghi</i> Brown Patch and Zoysia Patch <i>Rhizoctonia solani</i> Red Thread <i>Laetisaria fusiformis</i>	2 – 3-1/2	Apply when disease first appears. Make additional applications at 14-day intervals as needed. Allow spray to dry on leaves with no “watering-in”.
Pink Snow Mold <i>Microdochium nivale</i> (Only for those areas where snow cover is not present the entire winter)	2 – 3-1/2	Apply TRANSOM® 4.5 F in middle to late November before turf has stopped all growth activity. Lightly water application into the root zone for best results. For best results, use a spreader-sticker. Second spray should dry on leaf surfaces with no “watering-in”. Minimum spray interval is 14 days.
Gray Leaf Spot <i>Pyricularia grisea</i>	3-1/2 – 5-1/3	Apply when conditions are favorable for disease development. Continue applications at 14-day intervals. Allow spray to dry on leaves with no “watering-in”.
Summer Patch <i>Magnaportha poae</i>	3-1/2 – 5-1/3	For preventative treatment, make 3 applications at 21-day intervals beginning in early May. Water product into the root zone thoroughly after application. For suppression, apply two applications at 14-day intervals beginning when disease first appears.
Fusarium Blight <i>Fusarium</i> spp. Necrotic Ring Spot and Spring Dead Spot <i>Leptosphaeria korrae</i>	3-1/2 – 5-1/3	Make two applications at 14-day intervals beginning when disease first appears.
Stripe Smut <i>Ustilago striformis</i>	3-1/2 – 5-1/3	Make two applications at 14-day intervals when disease first appears. Water product into the root zone after application. For prevention, apply in the spring (just before grass begins growth), and in the fall.

*Use 3-1/2 to 5-1/3 fl. oz. rate for curative response to Basal Stem Anthracnose.

**Refer to the “Use Sites and Maximum Application Rates” table to determine allowable rates for each application.

Turf Application Directions

Apply TRANSOM® 4.5 F uniformly over the area to be treated with a properly calibrated power sprayer. Apply after mowing or avoid mowing for 12 hours after application. Apply sufficient water to obtain thorough coverage, usually 1-1/2 to 2-1/2 gallons per 1,000 sq. ft. of turf area. When treating golf greens, always treat aprons and approaches to golf greens. The minimum retreatment interval for turf applications is 14 days.

Use Sites and Maximum Application Rates

Site	Maximum Single Application Rate fl. oz./1,000 sq. ft.	Maximum Seasonal Application Rate fl. oz./1,000 sq. ft.
Golf Course Greens, Tees and Aprons	5-1/3	14-1/4
Golf Course Fairways (Except Florida)	3-1/2	3-1/2
Golf Course Fairways (Florida Only)	1-3/4	1-3/4
Residential and Public Areas (home lawns, parks, athletic fields, schools, day care centers)	1-3/4	7

Note to User: Do not graze animals on treated turf. Do not feed clippings to livestock or poultry. Shake well before using.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a dry, temperature-controlled area.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinseate is a violation of Federal Law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this Product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of PROKöZ, INC. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold PROKöZ, INC. and Seller harmless for any claims relating to such factors.

PROKöZ, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or PROKöZ, INC., and Buyer and User assume the risk of any such use. PROKöZ, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent permitted by law, neither PROKöZ, INC. nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF PROKöZ, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF PROKöZ, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

PROKöZ, INC. and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of PROKöZ, INC.

TRANSOM® is a trademark of PROKöZ, INC.

FUNGICIDE

TRANSOM® 4.5 F

Liquid Flowable Systemic Turf and Ornamental Fungicide

For control of a broad spectrum of diseases of bedding, flowering, herbaceous and tropical foliage plants, shrubs, trees and flowers, containerized woody shrubs and trees and turfgrass.

ACTIVE INGREDIENT:

Thiophanate-methyl..... 46.2%

OTHER INGREDIENTS:..... 53.8%**TOTAL:**..... 100.0%

(Contains 4.5 Pounds Thiophanate-methyl Per Gallon)

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

See attached label booklet for additional PRECAUTIONARY STATEMENTS and DIRECTIONS FOR USE.

EPA Reg. No. 72112-2

EPA Est. No. indicated by the first two letters of the batch number on this package.

(AL) 228-IL-002, (YU) 67545-AZ-001

Product of China or India.

Formulated in the United States with

U.S. and imported ingredients.

Manufactured for:
PROKōZ, Inc.
 1720 Windward Concourse
 Suite 150
 Alpharetta, GA 30005

Net Contents:

2.5 Gallon (9.46 liters)

EPA 20100222

PEEL HERE TO OPEN **13091000[1]**

03212013